



This is an official DHEC/CDC Health Advisory

Distributed via Health Alert Network
June 21, 2023, 4:30 PM
10557-DAD-06-21-2023-MEASLES

Guidance on Measles during the Summer Travel Season

Summary

As the summer travel season begins, the Centers for Disease Control and Prevention (CDC) is issuing this Health Alert Network (HAN) Health Advisory to remind clinicians and public health officials to provide guidance for measles prevention to international travelers and to be on alert for cases of [measles](#). Guidance is also provided for people planning international travel. Measles (rubeola) is extremely contagious; one person infected by measles can infect 9 out of 10 of their unvaccinated close contacts. As of [June 8, 2023](#), CDC has been notified of 16 confirmed U.S. cases of measles across 11 jurisdictions, with 14 (88%) linked to international travel. Based on current estimates, twice as many Americans are planning to travel internationally in 2023 compared with 2022. Many countries and popular travel destinations, such as [London](#), England, have experienced measles outbreaks in recent years. The United States has seen an [increase in measles cases](#) during the first 5 months of 2023, with 16 reported cases compared with 3 in 2022 during the same period. Most of these cases were among children who had not received measles-mumps-rubella (MMR) vaccine. To prevent measles infection and spread from importation, all U.S. residents should be up to date on their MMR vaccinations, especially prior to international travel regardless of the destination.

Background

[Measles](#) is a highly contagious viral illness that typically begins with a prodrome of fever, cough, coryza (runny nose), and conjunctivitis (pink eye), lasting 2-4 days prior to [rash](#) onset. Measles can cause severe health complications, including pneumonia, encephalitis (inflammation of the brain), and death. The virus is transmitted by direct contact with infectious droplets or by airborne spread when an infected person breathes, coughs, or sneezes. Measles virus can remain infectious in the air and on

surfaces for up to 2 hours after an infected person leaves an area. Measles is transmitted by contact with an infected person through coughing and sneezing. Infected people are contagious from 4 days before the rash starts through 4 days afterward. The incubation period for measles from exposure to fever is usually about 10 days (range 7 to 12 days), and from exposure to rash onset is usually about 14 days (range 7 to 21 days). [Declines in measles vaccination rates globally during the COVID-19 pandemic](#) have increased the risk of larger measles outbreaks worldwide, including in the United States.

Measles outbreaks are occurring in all [World Health Organization \(WHO\) regions](#). Large and disruptive outbreaks (≥ 20 reported measles cases per million population during a 12-month period) have been reported in the European, African, Eastern Mediterranean, Western Pacific, and Southeast Asian regions during 2023. In the United States, measles is commonly associated with unvaccinated U.S. travelers returning from other countries where measles is actively circulating. International visitors and returning U.S. travelers can expose U.S. residents in transit and after arrival, leading to additional cases and the possibility for larger outbreaks.

Recommendations for Healthcare Professionals

- Ensure that all patients without other evidence of immunity, especially those planning international travel, are up to date on [MMR vaccine](#) and [other recommended vaccines](#) before their international travel.
- [CDC recommends](#) that all U.S. residents older than age 6 months who will travel internationally, without evidence of immunity, receive MMR vaccine prior to departure.
 - Infants 6 through 11 months of age should receive one dose of MMR vaccine before departure. Infants who receive a dose of MMR vaccine before their first birthday should receive two more doses of MMR vaccine, the first of which should be administered when the child is 12 through 15 months of age and the second at least 28 days later.
 - Children 12 months of age or older should receive two doses of MMR vaccine, separated by at least 28 days.
 - Teenagers and adults without evidence of measles immunity should have documentation of two doses of MMR vaccine separated by at least 28 days.
- At least one of the following is considered evidence of measles immunity for international travelers: 1) birth before 1957, 2) documented administration of two doses of live measles virus vaccine (MMR, MMRV, or other measles-containing vaccine), or 3) laboratory (serologic) proof of immunity or laboratory confirmation of disease.
- Consider measles as a diagnosis in anyone with fever ($\geq 101^\circ\text{F}$ or 38.3°C) and a generalized maculopapular rash with cough, coryza, or conjunctivitis who has recently been abroad, especially in countries with ongoing [outbreaks](#) (e.g., India, Somalia, and Yemen).

- Be aware that some patients may develop a mild rash reaction in the three weeks following MMR vaccination, which does not typically require testing or public health intervention.

Recommendations for State and Local Health Departments

- State and local health departments have the lead in investigating measles cases and outbreaks.
- Measles is an immediately notifiable disease and should be reported promptly (within 24 hours) by the state health department to CDC (measlesreport@cdc.gov) and through the [National Notifiable Disease Surveillance System \(NNDSS\)](#).
- Establish measles case reporting within hospitals and to public health authorities.
- Record and report details about cases of measles, including adherence to recommended precautions and facility location(s) of index and secondary cases.
- If measles is identified, conduct active surveillance for any additional cases and facilitate transportation of specimens immediately to confirm diagnosis.

Recommendations for International Travelers

- Consult with your doctor several weeks before traveling abroad, regardless of your destination, to see if you or any dependents need MMR vaccine. You should be up-to-date on all routine vaccines, such as measles-mumps-rubella (MMR), tetanus, and flu. Depending on where you're going and what activities you plan, other vaccines may be recommended.
- Two doses of MMR vaccine provide better protection (97%) against measles than one dose (93%). Getting MMR vaccine is much safer than getting measles, mumps, or rubella.
- At least one of the following is considered evidence of measles immunity for international travelers: 1) birth before 1957, 2) documented administration of two doses of live measles virus vaccine (MMR, MMRV, or other measles-containing vaccine), or 3) laboratory (serologic) proof of immunity or laboratory confirmation of disease
- You should plan to be fully vaccinated (two doses) at least 2 weeks before you depart.
- After international travel, you should watch for signs and symptoms of measles for 3 weeks after you return to the United States. If you or your child gets sick with a rash and a high fever, call your doctor. Tell them you traveled abroad and whether you or your child have received MMR vaccine.
- Before your next trip, check your [destination](#) and CDC's [Global Measles Travel Health Notice](#) for more travel health advice, including where measles outbreaks have been reported.

Resources for Additional Information

- International Travelers
 - [Plan for Travel - Measles | CDC](#)
 - [Safety Information for Measles, Mumps, Rubella \(MMR\) Vaccines | Vaccine Safety | CDC](#)
- Healthcare and Public Health Professionals
 - [For Healthcare Professionals - Diagnosing and Treating Measles | CDC](#)
 - [Interim Measles Infection Prevention Recommendations in Healthcare Settings | CDC](#)
 - [Measles - Vaccine Preventable Diseases Surveillance Manual | CDC](#)
 - [Rubeola / Measles | CDC Yellow Book 2024](#)

DHEC contact information for reportable diseases and reporting requirements

Reporting of **Measles** is consistent with South Carolina Law requiring the reporting of diseases and conditions to your state or local public health department. (State Law # 44-29-10 and Regulation # 61-20) as per the DHEC 2023 List of Reportable Conditions available at: <https://www.scdhec.gov/sites/default/files/Library/CR-009025.pdf>

Federal HIPAA legislation allows disclosure of protected health information, without consent of the individual, to public health authorities to collect and receive such information for the purpose of preventing or controlling disease. (HIPAA 45 CFR §164.512).

Regional Public Health Offices – 2023

Mail or call reports to the Epidemiology Office in each Public Health Region

MAIL TO:

<p style="text-align: center;">Lowcountry</p> <p>4050 Bridge View Drive, Suite 600 N. Charleston, SC 29405 Fax: (843) 953-0051</p>	<p style="text-align: center;">Midlands</p> <p>2000 Hampton Street Columbia, SC 29204 Fax: (803) 576-2993</p>	<p style="text-align: center;">Pee Dee</p> <p>1931 Industrial Park Road Conway, SC 29526 Fax: (843) 915-6506</p>	<p style="text-align: center;">Upstate</p> <p>352 Halton Road Greenville, SC 29607 Fax: (864) 282-4373</p>
---	--	---	---

CALL TO:

<p style="text-align: center;">Lowcountry</p> <p>Allendale, Bamberg, Beaufort, Berkeley, Calhoun, Charleston, Colleton, Dorchester, Hampton, Jasper, Orangeburg</p> <p>Office: (843) 441-1091 Nights/Weekends: (843) 441-1091</p>	<p style="text-align: center;">Midlands</p> <p>Aiken, Barnwell, Chester, Edgefield, Fairfield, Kershaw, Lancaster, Lexington, Newberry, Richland, Saluda, York</p> <p>Office: (888) 801-1046 Nights/Weekends: (888) 801-1046</p>	<p style="text-align: center;">Pee Dee</p> <p>Clarendon, Chesterfield, Darlington, Dillon, Florence, Georgetown, Horry, Lee, Marion, Marlboro, Sumter, Williamsburg</p> <p>Office: (843) 915-8886 Nights/Weekends: (843) 409-0695</p>	<p style="text-align: center;">Upstate</p> <p>Abbeville, Anderson, Cherokee, Greenville, Greenwood, Laurens, McCormick, Oconee, Pickens, Spartanburg, Union</p> <p>Office: (864) 372-3133 Nights/Weekends: (864) 423-6648</p>
--	---	--	--

<p style="text-align: center;">For information on reportable conditions, see https://www.scdhec.gov/ReportableConditions</p>	<p><u>DHEC Bureau of Communicable Disease Prevention & Control</u> Division of Acute Disease Epidemiology 2100 Bull St • Columbia, SC 29201 Phone: (803) 898-0861 • Fax: (803) 898-0897 Nights / Weekends: 1-888-847-0902</p>
---	---

Categories of Health Alert messages:

Health Alert	Conveys the highest level of importance; warrants immediate action or attention.
Health Advisory	Provides important information for a specific incident or situation; may not require immediate action.
Health Update	Provides updated information regarding an incident or situation; unlikely to require immediate action.
Info Service	Provides general information that is not necessarily considered to be of an emergent nature.