

South Carolina Department of Health and Environmental Control Division of Acute Disease Epidemiology

Week Ending October 24, 2015 (MMWR Week 42)

All data are provisional and may change as more reports are received.

In this issue:	
Summary	2
I. Confirmatory testing	3
II. Positive rapid tests	6
III. ILINet	7
IV. Hospitalizations and deaths	10
V. National influenza	12
surveillance	
VI. SC influenza surveillance	13
components	
VII. Definitions for influenza surveillance	14

MMWR Week 42 at a Glance:

Influenza Activity Synopsis:

During MMWR week 42 influenza activity in South Carolina remained fairly low. South Carolina reported LOCAL activity.

Laboratory surveillance:

- 118 laboratory-confirmed cases of influenza were reported from 20 counties.
- 176 (56.6%) of all laboratory-confirmed cases this season are influenza A, 124 (39.9%) are influenza B, 5 (1.6%) are influenza A/B, and 6 (1.9%) is influenza unknown subtype.

ILI Activity (South Carolina baseline is 2.05%):

The percentage of visits to sentinel providers for influenzalike illness (1.87%) was below South Carolina's baseline.
 ILI percentages represent ILI activity reported by less than half of enrolled sentinel providers. Therefore, ILI percentages may not be representative of actual flu activity.

Hospitalizations:

• 20 laboratory confirmed influenza-associated hospitalizations were reported. Since October 4, 2015, 50 laboratory confirmed influenza associated hospitalizations have been reported.

Deaths:

 No laboratory confirmed influenza-associated deaths were reported. Since October 4, 2015 five laboratory confirmed influenza associated deaths have been reported.

Summary of Laboratory Confirmed Tests, ILI Activity, Influenza Associated Hospitalizations and Deaths Compared to Previous Week and Previous Season

	Current week	Previous week	Change from previous week	Cumulative (2015-16)	Cumulative (2014-15)	Cumulative change 2015-16 compared to 2014-15
Number of positive confirmatory tests (culture, RT-PCR, DFA, IFA)	1	0	A	2	5	V 60.0%
Positive rapid antigen tests	117	114	A 2.6%	309	318	lacksquare
Percent of ILI visits reported by ILINet providers	1.87%	3.16%	V 1.29%			
Number of lab confirmed flu hospitalizations	20	19	▲ 5.3%	50	27	▲ 85.2%
Number of lab confirmed flu deaths	0	1	V 75.0%	5	4	\$\times 25.0%


I. Confirmatory testing

Positive confirmatory influenza test results* Current MMWR Week (10/18/15 – 10/24/15)						
	BOL and reference labs					
Number of positive confirmatory tests 1						
Influenza A unsubtyped	0					
Influenza A H1N1 0						
Influenza A H3N2 1						
Influenza B	0					
Other	0					
Includes culture, RT-PCR, DFA, and IFA						

For the current MMWR reporting week, 1 positive confirmatory test was reported.


Positive confirmatory influenza test results* Cumulative (10/4/15 – 10/24/15)						
	BOL and reference labs					
Number of positive confirmatory tests 2						
Influenza A unsubtyped	1					
Influenza A H1N1 0						
Influenza A H3N2 1						
Influenza B	0					
Other 0						
Includes culture, RT-PCR, DFA, and IFA						

Positive Confirmatory Tests by County* Current Week 10/18/15 – 10/24/15



^{*}Includes culture, PCR, DFA, IFA


Map of all Laboratory Confirmed Cases (n)* and Population Case Rates/100,000 by County Cumulative 10/4/15 – 10/24/15





II. Positive Rapid Antigen Tests

For the current MMWR reporting week, 117 positive rapid antigen tests were reported. Of these, 66 were influenza A, 50 were influenza B, and 1 was influenza AB. This compares to 144 during this same week last year.

Map of Positive Rapid Influenza Tests by County (10/18/15 - 10/24/15)








III. ILINet Influenza-Like Illness Surveillance

During the most recent MMWR week, 1.87%* of patient visits to SC ILINet providers were due to ILI. This is below the state baseline (2.05%). This ILI percentage compares to 1.92% this time last year. Reports were received from providers in 10 counties, representing all of the 4 regions.



^{*}The SC baseline is the mean percentage of patient visits for ILI during non-influenza weeks (weeks when percent of positive lab tests was below 20%) for the previous three seasons plus two standard deviations. * ILI percentage is dependent upon the number of reporting providers and can be greatly influenced by a single provider with high numbers of ILI.

Influenza-Like Illness Reported by Sentinel Providers October 18, 2015 – October 24, 2015

County	ILI %	County	ILI %	
Abbeville		Greenwood	NR	
Aiken	0%	Hampton	NR	
Allendale		Horry		
Anderson	1.59%	Jasper		
Bamberg		Kershaw		
Barnwell		Lancaster		
Beaufort	NR	Laurens	NR	
Berkeley	3.70%	Lee		
Calhoun		Lexington	NR	
Charleston	3.99%	Marion		
Cherokee		Marlboro		
Chester		McCormick	NR	
Chesterfield		Newberry		
Clarendon		Oconee		
Colleton		Orangeburg		
Darlington		Pickens	0%	
Dillon	<u></u>	Richland	0%	
Dorchester	NR	Saluda	0%	
Edgefield		Spartanburg	NR	
Fairfield		Sumter	NR	
Florence	0.57%	Union		
Georgetown	NR	Williamsburg		
Greenville	0%	York	0%	

NR: No reports received ---: No enrolled providers


IV. Influenza-associated hospitalizations and deaths


For the current MMWR reporting week, 20 laboratory confirmed influenza-associated hospitalizations were reported by 49 hospitals. No laboratory confirmed influenza-associated deaths were reported. Since October 4, 2015 50 laboratory confirmed influenza-associated hospitalizations and 5 laboratory confirmed influenza-associated deaths have been reported. Laboratory confirmation for hospitalizations and deaths includes culture, PCR, DFA,

Current MMWR Week (10/18/15 - 10/24/15)							
	0-4	5-17	18-49	50-64	65+	Unknown	Total
Hospitalizations	2	1	2	6	9	0	20
Deaths	0	0	0	0	0	0	0

Cumulative (10/4/15 -10/24/14)							
	0-4	5-17	18-49	50-64	65+	Unknown	Total
Hospitalizations	5	1	5	12	27		50
Deaths	0	0	1	1	3		5

^{*} Lab confirmation for hospitalizations and deaths includes culture, PCR, DFA, IFA, and rapid test.






V. National surveillance (10/11/15 - 10/17/15)

During week 41 (October 11-17, 2015), influenza activity was low in the United States.

- Viral Surveillance: The most frequently identified influenza virus type reported by public health laboratories in week 41 was influenza A viruses, with influenza A (H3) viruses predominating. The percentage of respiratory specimens testing positive for influenza in clinical laboratories is low.
- Pneumonia and Influenza Mortality: The proportion of deaths attributed to pneumonia and influenza (P&I) was below their system-specific epidemic threshold in both the NCHS Mortality Surveillance System and the 122 Cities Mortality Reporting System.
- o <u>Influenza-associated Pediatric Deaths:</u> No influenza-associated pediatric deaths were reported.
- Outpatient Illness Surveillance: The proportion of outpatient visits for influenza-like illness (ILI) was 1.4%, which is below the national baseline of 2.1%. All 10 regions reported ILI below region-specific baseline levels. Puerto Rico, New York City and 50 states experienced minimal ILI activity and the District of Columbia had insufficient data.
- Geographic Spread of Influenza: The geographic spread of influenza in Guam was reported as widespread; four states reported local activity; Puerto Rico and 29 states reported sporadic activity; and the District of Columbia, the U.S. Virgin Islands and 17 states reported no influenza activity.

Influenza-Like Illness (ILI) Activity Level Indicator Determined by Data Reported to ILINet 2015-16 Influenza Season Week 41 ending Oct 17, 2015



VI. South Carolina Influenza Surveillance Components

South Carolina influenza surveillance consists of mandatory and voluntary reporting systems for year-round influenza surveillance. These networks provide information on influenza virus strain and subtype and influenza disease burden.

Mandatory Reporting

Positive confirmatory test reporting

Positive influenza culture, PCR, DFA, and IFA results from commercial laboratories must be reported to DHEC within 3 days electronically via CHESS or using a DHEC 1129 card.

Positive rapid antigen test reporting

Summary numbers of positive rapid influenza tests and influenza type identified must be sent to the regional health department by fax or email before noon on Monday for the preceding week.

Influenza deaths

All laboratory confirmed influenza deaths (adult and pediatric) must be reported to DHEC within 24 hours. These include results from viral culture, PCR, rapid flu tests, DFA, IFA or autopsy results consistent with influenza.

Influenza hospitalizations

DHEC requires weekly submission of laboratory confirmed influenza hospitalizations. Hospitals must report these to their <u>regional</u> health department by noon on Monday for the preceding week.

For additional information about ILINet or to become an ILINet provider, contact the Acute Disease Epidemiology influenza surveillance coordinator at springcb@dhec.sc.gov.

Voluntary Networks

Influenza-Like Illness (ILINet) Sentinel Providers Network ILINet focuses on the number of patients presenting with influenza-like symptoms in the absence of another known cause. ILI is defined as fever (temperature ≥100°F) plus a cough and/or a sore throat in the absence of another known cause. Providers submit weekly reports to the CDC of the total number of patients seen in a week and the subset number of those patients with ILI symptoms by age group.

South Carolina Disease Alerting, Reporting & Tracking System (SC-DARTS)

SC-DARTS is a collaborative network of syndromic surveillance systems within South Carolina. Currently our network contains the following data sources: SC Hospital Emergency Department (ED) chief-complaint data, Poison Control Center call data, Over-the-Counter (OTC) pharmaceutical sales surveillance, and CDC's BioSense Biosurveillance system. The hospital ED syndromic surveillance system classifies ED chief complaint data into appropriate syndrome categories (ex: Respiratory, GI, Fever, etc.). These syndrome categories are then analyzed using the cumulative sum (CUSUM) methodology to detect any significant increases. Syndromic reports are distributed back to the hospital on a daily basis.

VII. Definitions for Influenza Surveillance

Activity level: Indicator of the geographic spread of influenza activity which is reported to CDC each week.

- No activity: No increase in ILI activity and no laboratory-confirmed influenza cases.
- Sporadic: No increase in ILI activity and isolated laboratory-confirmed influenza cases
- *Local:* Increased ILI or 2 or more institutional outbreaks in one region and laboratory-confirmed influenza cases within the past 3 weeks in the region with increased ILI or outbreaks
- **Regional:** Increased ILI or institutional outbreaks in 2-3 regions and laboratory-confirmed influenza cases within the past 3 weeks in the regions with increased ILI or institutional outbreaks
- *Widespread:* Increased ILI and/or institutional outbreaks in at least 4 regions and laboratory confirmed influenza in the state within the past 3 weeks

Confirmatory testing: Influenza testing which is considered to be confirmatory, such as a viral culture or RT-PCR

Influenza-associated death: A death in which laboratory confirmation (see definition below) for influenza was reported, or for which an autopsy report consistent with influenza was provided, regardless of primary cause of death.

Influenza-like illness (ILI): Fever (temperature of 100°F [37.8°C] or greater) and cough and/or sore throat

MMWR week: Term for influenza surveillance week. Each week begins on Sunday and ends on Monday. The influenza season begins with MMWR week 40 and ends with MMWR week 39. The 2015-16 influenza season began on October 4, 2015 and will end on October 1, 2016.

Laboratory-confirmation: Influenza positive resulting from one of the following methods:

- DFA
- IFA
- Rapid influenza antigen test
- RT-PCR
- Viral culture