Hospital Infections Disclosure Act Report

Reported by: South Carolina Department of Health and Environmental Control

Surgical Site Infection (SSI) Standardized Infection Ratio by Procedure

Data Collected: 01/01/2017 - 06/30/2017

Procedure	No. of Specific Procedures Performed ^a	No. of Infections	No. of Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval
Coronary Bypass Graft (Chest and Donor Incision)	154	0	1.04	0.00	, 2.873
Coronary Bypass Graft (Chest Only Incision)	14	*	*	*	*
Colon Surgery	151	3	4.19	0.72	0.182, 1.947
Hip Prosthesis (Replacement)	141	2	1.02	1.95	0.327, 6.453
Abdominal Hysterectomy	78	0	0.48	*	*
Knee Prosthesis (Replacement)	301	3	0.91	*	*

a. *= Too few procedures. Reporting on too few procedures is a risk to patient confidentiality and data stability. If less than twenty surgical procedures are performed, the SIR and number of infections will be suppressed until more procedures are performed.

Central Line Associated Blood Stream Infection (CLABSI) Standardized Infection Ratio (SIR)

Data Collected: 01/01/2017 - 06/30/2017

Location ^a	No. of Central Line Days ^{b,c}	No. of Infections	No. of Predicted Infections	Standardized Infection Ratio	95% Confidence Interval
All Adult Critical Care Units	7226	4	8.03	0.5	0.136,1.275
All Adult Inpatient Wards	8986	9	7.72	1.2	0.533,2.213
All Pediatric Critical Care Units	148	0	0.19	*	*
All Pediatric Inpatient Wards	28	*	0.02	*	*
Adult Speciality Care	3962	0	4.21	0.0	0,0.876
Neonatal Intensive Care Unit	430	0	0.78	*	*

a. The specific patient care area in which a patient is assigned while receiving care in the healthcare facility. All adult and pediatric critical care units are combined into one SIR; all adult and pediatric inpatient wards are combined into one SIR for this report.

b. Central line days are the total number of days a central line is in place for all patients in selected hospital locations.

c. * = Too few central line days. Reporting on too few central line days is a risk to confidentiality and data stability. If there are less than fifty central line days, the SIR and number of infections will be suppressed until there are more central line days to report.

Methicillin-resistant Staphylococcus aureus bloodstream infection (MRSA BSI) LabID Event Data Facility Wide Inpatient Data Collected: 01/01/2017 - 06/30/2017

Hospital Onset MRSA BSI Standardized Infection Ratio (SIR)				
No. Patient Days	No. Patient Days No. LabID Events a Predicted No. of LabID Events SIR			95% Confidence Interval
73722	10	6.2041107	1.612	0.819, 2.873

a. Hospital Onset: LabID event specimen collected as an inpatient >3 days after admission to the facility (i.e., on or after day 4)

Clostridium Difficile Infections(CDI) LabID Event Data

Facility Wide Inpatient Data Collected: 01/01/2017 - 06/30/2017

Hospital Onset CDI LabID Event Data					
No. Patient Days No. of LabID Events a Predicted No. of LabID Events SIR Confident Confidence of LabID Events SIR Confidenc				95% Confidence Interval	
66097	39	50.445783	0.773	0.557, 1.046	

a. Hospital Onset: LabID event specimen collected as an inpatient >3 days after admission to the facility (i.e., on or after day 4)

Ventilator Associated Events(VAE) Rate

Data Collected: 01/01/2017 - 06/30/2017

Ventilator Associated Events(VAE)				
No. Ventilator Days No. Ventilator Days No. Predicted No. of IVAC-plus Events a of IVAC-Plus Events		SIR	95% Confidence Interval	
4203	20	14.125533	1.416	0.889, 2.148

a. IVAC-plus Events: All Ventilator associated events meeting the Infection-related Ventilator Associated Complications (IVAC) and Possible Ventilator-associated pneumonia (PVAP) definitions